

Göteborg ITP 98-07
(Revised version)

Open groups of constraints - Integrating arbitrary involutions -

Igor Batalin¹ and Robert Marnelius²

*Institute of Theoretical Physics
Chalmers University of Technology
Göteborg University
S-412 96 Göteborg, Sweden*

Abstract

A new type of quantum master equation is presented which is expressed in terms of a recently introduced quantum antibracket. The equation involves only two operators: an extended nilpotent BFV-BRST charge and an extended ghost charge. It is proposed to determine the generalized quantum Maurer-Cartan equations for arbitrary open groups. These groups are the integration of constraints in arbitrary involutions. The only condition for this is that the constraint operators may be embedded in an odd nilpotent operator, the BFV-BRST charge. The proposal is verified at the quasigroup level. The integration formulas are also used to construct a generating operator for quantum antibrackets of operators in arbitrary involutions.

¹On leave of absence from P.N.Lebedev Physical Institute, 117924 Moscow, Russia
E-mail: batalin@td.lpi.ac.ru.

²E-mail: tferm@fy.chalmers.se

1 Introduction.

In [1] we have introduced quantum antibrackets. They are new objects which cast new light on the BV-quantization of arbitrary gauge theories. Here we show that they are also useful in a more general context. In terms of the quantum antibracket we give a new simple quantum master equation which we conjecture to encode the Maurer-Cartan equations and their generalizations to open groups whose generators are in arbitrary involutions. This conjecture is verified at the quasigroup level. The new master equation involves only two operators: an extended nilpotent BFV-BRST charge and an extended ghost charge. The construction is thus embedded in a general BRST framework which should make the formulas useful for general gauge theories.

In section 2 we review the BFV-treatment of constraints in arbitrary involutions, and in section 3 we present the new master equation for generalized quantum Maurer-Cartan equations. In section 4 we treat quasigroup theories in detail, and in section 5 the formalism is used to generalize the generating operator for quantum antibrackets given in [1] to operators in arbitrary involutions.

2 Constraints in arbitrary involutions.

Consider classical, real constraints θ_a on a symplectic manifold. Their Grassmann parities are $\varepsilon(\theta_a) \equiv \varepsilon_a (= 0, 1)$. θ_a are assumed to be in arbitrary involutions with respect to the Poisson bracket, *i.e.*

$$\{\theta_a, \theta_b\} = U_{ab}{}^c \theta_c, \quad (1)$$

where the structure coefficients $U_{ab}{}^c$ may be arbitrary functions on the considered symplectic manifold. It is well-known that this algebra always may be embedded in one single real, odd function Ω on a ghost extended manifold in such a way that $\{\Omega, \Omega\} = 0$ in terms of the extended Poisson bracket. Ω is the BFV-BRST charge [2]. The corresponding quantum theory is consistent if the corresponding odd, hermitian operator Ω is nilpotent, *i.e.* $\Omega^2 = 0$. For a finite number of degrees of freedom such a solution always exists and is of the form [3] (N is the rank of the theory and θ_a are the hermitian constraint operators)

$$\Omega = \sum_{i=0}^N \Omega_i, \quad (2)$$

$$\Omega_0 \equiv \mathcal{C}^a \theta_a, \quad \Omega_i \equiv \Omega_{a_1 \dots a_{i+1}}^{b_1 \dots b_{i+1}} (\mathcal{P}_{b_1} \dots \mathcal{P}_{b_i} \mathcal{C}^{a_{i+1}} \dots \mathcal{C}^{a_1})_{Weyl}, \quad i = 1, \dots, N, \quad (3)$$

where we have introduced the ghost operators $\mathcal{C}^a$, $\mathcal{P}_a$, $\varepsilon(\mathcal{C}^a) = \varepsilon_a + 1$, satisfying

$$[\mathcal{C}^a, \mathcal{P}_b] = i\hbar \delta_b^a, \quad (\mathcal{C}^a)^\dagger = \mathcal{C}^a, \quad \mathcal{P}_a^\dagger = -(-1)^{\varepsilon_a} \mathcal{P}_a. \quad (4)$$

In (3) the ghost operators are Weyl ordered which means that Ω_i are all hermitian. One may notice that Ω in (2) has ghost number one, *i.e.*

$$[G, \Omega] = i\hbar \Omega, \quad G \equiv -\frac{1}{2} (\mathcal{P}_a \mathcal{C}^a - \mathcal{C}^a \mathcal{P}_a (-1)^{\varepsilon_a}), \quad (5)$$

where G is the ghost charge. Ω determines the precise form of the quantum counterpart of the algebra (1). A convenient form of this algebra is obtained if we rewrite Ω in the following $\mathcal{CP}$ -ordered form [3]

$$\begin{aligned}\Omega &= \sum_{i=0}^N \Omega'_i, \quad \Omega'_0 \equiv \mathcal{C}^a \theta'_a, \\ \Omega'_i &\equiv \mathcal{C}^{a_{i+1}} \dots \mathcal{C}^{a_1} \Omega'_{a_1 \dots a_{i+1}}{}^{b_1 \dots b_i} \mathcal{P}_{b_1} \dots \mathcal{P}_{b_i}, \quad i = 1, \dots, N.\end{aligned}\tag{6}$$

The nilpotency of Ω requires then the algebra

$$[\theta'_a, \theta'_b] = i\hbar U'_{ab}{}^c \theta'_c,\tag{7}$$

where the structure operators $U'_{ab}{}^c$ are given by

$$U'_{ab}{}^c = 2(-1)^{\varepsilon_b + \varepsilon_c} \Omega'_{ab}{}^c.\tag{8}$$

In terms of the coefficient operators in (3) $\Omega'_{ab}{}^c$ and θ'_a are given by

$$\begin{aligned}\Omega'_{ab}{}^c &= \Omega_{ab}^c + \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{i\hbar}{2} \right)^n (n+1)(n+2)! \Omega_{aba_1 \dots a_n}^{a_n \dots a_1 c} (-1)^{\sum_{k=1}^n \varepsilon_{a_k}}, \\ \theta'_a &= \theta_a + \sum_{n=1}^{\infty} \left(\frac{i\hbar}{2} \right)^n (n+1)! \Omega_{aa_1 \dots a_n}^{a_n \dots a_1} (-1)^{\sum_{k=1}^n \varepsilon_{a_k}},\end{aligned}\tag{9}$$

which shows that θ'_a in general are different from θ_a and in general not even hermitian. However, the main point here is that Ω through *e.g.* (7) represents the quantum counterpart of (1).

3 Quantum master equation and generalized Maurer-Cartan equations.

We want now to integrate the quantum involution (1) encoded in Ω as represented by *e.g.* (7), or in other words we are looking for finite gauge transformations if θ'_a are viewed as gauge generators. We consider therefore the Lie equations

$$A(\phi) \overleftarrow{\nabla}_a \equiv A(\phi) \overleftarrow{\partial}_a - (i\hbar)^{-1} [A(\phi), Y_a(\phi)] = 0,\tag{10}$$

where ∂_a is a derivative with respect to the parameter ϕ^a , $\varepsilon(\phi^a) = \varepsilon_a$. The operator Y_a , which depends on ϕ^a , must satisfy integrability conditions

$$Y_a \overleftarrow{\partial}_b - Y_b \overleftarrow{\partial}_a (-1)^{\varepsilon_a \varepsilon_b} = (i\hbar)^{-1} [Y_a, Y_b],\tag{11}$$

which in turn are integrable without further conditions. In order for the Lie equations (10) to be connected to the integration of the quantum involution (1) $Y_a(\phi)$ has to be of the form

$$Y_a(\phi) = \lambda_a^b(\phi) \theta'_b (-1)^{\varepsilon_a + \varepsilon_b} + \{\text{possible ghost dependent terms}\}, \quad \lambda_a^b(0) = \delta_a^b,\tag{12}$$

where $\lambda_a^b(\phi)$ are operators in general. One may note that in a ghost independent scheme θ'_a are the generators of the finite transformations. However, within our BRST framework,

$[\Omega, \mathcal{P}_a] = \theta'_a + \{\text{ghost dependent terms}\}$, are the appropriate generators. This motivates the form (12) and it also makes it natural to expect Y_a to have the general form

$$Y_a(\phi) = (i\hbar)^{-1}[\Omega, \Omega_a(\phi)], \quad \varepsilon(\Omega_a) = \varepsilon_a + 1, \quad (13)$$

where Y_a has ghost number zero and Ω_a ghost number minus one. This form implies that $[Y_a(\phi), \Omega] = 0$, so that if $[A(0), \Omega] = 0$ then $[A(\phi), \Omega] = 0$, *i.e.* a BRST invariant operator remains BRST invariant when transformed according to (10). In the following we consider Y_a to be given by Ω_a through equation (13). Equations (12) and (13) together with (2) and (3) imply that Ω_a in (13) must be of the form

$$\Omega_a(\phi) = \lambda_a^b(\phi)\mathcal{P}_b + \{\text{possible ghost dependent terms}\}, \quad \lambda_a^b(0) = \delta_a^b. \quad (14)$$

The integrability condition (11) for Y_a leads then by means of (13) to the following equivalent equation for Ω_a

$$\Omega_a \overleftarrow{\partial}_b - \Omega_b \overleftarrow{\partial}_a (-1)^{\varepsilon_a \varepsilon_b} = (i\hbar)^{-2}(\Omega_a, \Omega_b)_\Omega - \frac{1}{2}(i\hbar)^{-1}[\Omega_{ab}, \Omega], \quad (15)$$

where we have introduced the quantum antibracket defined in accordance with [1], *i.e.*

$$\begin{aligned} (A, B)_Q &\equiv \frac{1}{2} \left([A, [Q, B]] - [B, [Q, A]](-1)^{(\varepsilon_A+1)(\varepsilon_B+1)} \right) = \\ &= \frac{1}{2} \left([[A, Q], B] - [[B, Q], A](-1)^{(\varepsilon_A+1)(\varepsilon_B+1)} \right), \quad Q^2 = 0. \end{aligned} \quad (16)$$

This expression satisfies all properties of an antibracket if A and B commute, or for more general A 's and B 's under certain conditions [1]. Due to the form (14) of Ω_a , eq.(15) are generalized Maurer-Cartan equations for $\lambda_a^b(\phi)$. The integrability conditions of (15) lead to equivalent first order equations of Ω_{ab} and so on. Thus, Y_a is replaced by a whole set of operators, and the integrability condition (11) for Y_a is replaced by a whole set of integrability conditions. What is amazing is that (15) together with all its integrability conditions seem to be possible to embed in one single quantum master equation (eq.(20) below) which involves only two operators. The first operator is an extended nilpotent BFV-BRST charge involving the conjugate momenta π_a to ϕ^a , now turned into operators, and new ghost variables η^a , $\varepsilon(\eta^a) = \varepsilon_a + 1$, to be treated as parameters. It is

$$\Delta \equiv \Omega + \eta^a \pi_a (-1)^{\varepsilon_a}, \quad \Delta^2 = 0, \quad [\phi^a, \pi_b] = i\hbar \delta_b^a. \quad (17)$$

The second operator in the master equation is an even operator $S(\phi, \eta)$ defined by

$$\begin{aligned} S(\phi, \eta) &\equiv G + \eta^a \Omega_a(\phi) + \frac{1}{2} \eta^b \eta^a \Omega_{ab}(\phi) (-1)^{\varepsilon_b} + \\ &+ \frac{1}{6} \eta^c \eta^b \eta^a \Omega_{abc}(\phi) (-1)^{\varepsilon_b + \varepsilon_a \varepsilon_c} + \dots \\ &\dots + \frac{1}{n!} \eta^{a_n} \dots \eta^{a_1} \Omega_{a_1 \dots a_n}(\phi) (-1)^{(\varepsilon_{a_2} + \dots + \varepsilon_{a_{n-1}} + \varepsilon_{a_1} \varepsilon_{a_n})} + \dots, \end{aligned} \quad (18)$$

where G is the ghost charge operator in (5). Our main conjecture is that the operators $\Omega_{a_1 \dots a_n}(\phi)$ in (18) may be identified with Ω_a, Ω_{ab} in (15) and all the Ω 's in their integrability conditions in a particular manner. They satisfy the properties

$$\varepsilon(\Omega_{a_1 \dots a_n}) = \varepsilon_{a_1} + \dots + \varepsilon_{a_n} + n, \quad [G, \Omega_{a_1 \dots a_n}] = -ni\hbar \Omega_{a_1 \dots a_n}. \quad (19)$$

The last relation implies that $\Omega_{a_1 \dots a_n}$ has ghost number minus n . If we assign ghost number one to η^a then Δ has ghost number one and S has ghost number zero. The operators $\Omega_{a_1 \dots a_n}(\phi)$ are determined by the following quantum master equation

$$(S, S)_\Delta = i\hbar[\Delta, S], \quad (20)$$

where $(S, S)_\Delta$ is the quantum antibracket defined in accordance with (16). Thus, we have

$$(S, S)_\Delta \equiv [[S, \Delta], S]. \quad (21)$$

Consistency requires $[\Delta, S]$ to be nilpotent since

$$[\Delta, (S, S)_\Delta] = 0 \Leftrightarrow [\Delta, S]^2 = 0. \quad (22)$$

The explicit form of $[S, \Delta]$ to the lowest orders in η^a is

$$\begin{aligned} [S, \Delta] &= i\hbar\Omega + \eta^a[\Omega_a, \Omega] + \eta^b\eta^a\Omega_a \overleftarrow{\partial}_b i\hbar(-1)^{\varepsilon_b} + \frac{1}{2}\eta^b\eta^a[\Omega_{ab}, \Omega](-1)^{\varepsilon_b} + \\ &+ \frac{1}{2}\eta^c\eta^b\eta^a\Omega_{ab} \overleftarrow{\partial}_c i\hbar(-1)^{\varepsilon_b+\varepsilon_c} + \frac{1}{6}\eta^c\eta^b\eta^a[\Omega_{abc}, \Omega](-1)^{\varepsilon_b+\varepsilon_a\varepsilon_c} + O(\eta^4). \end{aligned} \quad (23)$$

To zeroth and first order in η^a the master equation (20) is satisfied identically. However, to second order in η^a it yields exactly (15). At third order in η^a it yields

$$\begin{aligned} \partial_a\Omega_{bc}(-1)^{\varepsilon_a\varepsilon_c} + \frac{1}{2}(i\hbar)^{-2}(\Omega_a, \Omega_{bc})_\Omega(-1)^{\varepsilon_a\varepsilon_c} + \text{cycle}(a, b, c) = \\ = -(i\hbar)^{-3}(\Omega_a, \Omega_b, \Omega_c)_\Omega(-1)^{\varepsilon_a\varepsilon_c} - \frac{2}{3}(i\hbar)^{-1}[\Omega'_{abc}, \Omega], \\ \Omega'_{abc} \equiv \Omega_{abc} - \frac{1}{8} \left\{ (i\hbar)^{-1}[\Omega_{ab}, \Omega_c](-1)^{\varepsilon_a\varepsilon_c} + \text{cycle}(a, b, c) \right\}, \end{aligned} \quad (24)$$

where we have introduced a higher quantum antibracket defined by (this expression may be obtained from appendix B in [1])

$$\begin{aligned} (A, B, C)_Q(-1)^{(\varepsilon_A+1)(\varepsilon_C+1)} &\equiv \\ &\equiv \frac{1}{3} \left([(A, B)_Q, C](-1)^{\varepsilon_C+(\varepsilon_A+1)(\varepsilon_C+1)} + \text{cycle}(A, B, C) \right), \quad Q^2 = 0, \end{aligned} \quad (25)$$

where the quantum antibracket on the right-hand side is given by (16). It satisfies Leibniz' rule if (16) does. The quantum antibracket (16) satisfies the Jacobi identities only if $(A, B, C)_Q = 0$ since we have

$$\begin{aligned} (A, (B, C)_Q)_Q(-1)^{(\varepsilon_A+1)(\varepsilon_C+1)} + \text{cycle}(A, B, C) = \\ = -\frac{1}{2}[(A, B, C)_Q(-1)^{(\varepsilon_A+1)(\varepsilon_C+1)}, Q] \end{aligned} \quad (26)$$

(cf. eq.(36) in [1]).

Comparing equation (24) and the integrability conditions of (15) we find exact agreement. We have also checked that the consistency condition (22) yields exactly (11) to second order in η^a , which is consistent with (15) as it should. Similarly we have checked that (22) to third order in η^a yields a condition which is consistent with (24), exactly like (11) is consistent with (15).

The master equation (20) yields at higher orders in η^a equations involving still higher quantum Ω -antibrackets defined in terms of lower antibrackets like in (25), and operators $\Omega_{abc\dots}$ with still more indices. We conjecture that these equations agree exactly with the integrability conditions of (24). For a rank- N theory we expect that there exists a solution of the form (18) to the master equation (20), which terminates just at the maximal order η^N (for a particular example see next section).

4 Example: Quasigroup first rank theories.

As an illustration of our formulas we consider now constraint operators θ_a forming a rank one theory in which case we have

$$\Omega = \mathcal{C}^a \theta'_a + \frac{1}{2} \mathcal{C}^b \mathcal{C}^a U_{ab}^c \mathcal{P}_c (-1)^{\varepsilon_c + \varepsilon_b}, \quad \theta'_a \equiv \theta_a + \frac{1}{2} i\hbar U_{ab}^b (-1)^{\varepsilon_b}. \quad (27)$$

The nilpotence of Ω requires (7) and

$$\left(i\hbar U_{ab}^d U_{dc}^e + [U_{ab}^e, \theta'_c] (-1)^{\varepsilon_c \varepsilon_e} \right) (-1)^{\varepsilon_a \varepsilon_c} + \text{cycle}(a, b, c) \equiv 0, \quad (28)$$

and in addition also restrictions on the commutators $[U_{ab}^c, U_{de}^f]$. The latter condition is satisfied if

$$[U_{ab}^c, U_{de}^f] = 0, \quad [[\theta_d, U_{ab}^c], U_{ef}^g] = 0, \quad (29)$$

which corresponds to quasigroups [4]. In this case Ω_a may be chosen to be

$$\Omega_a(\phi) = \lambda_a^b(\phi) \mathcal{P}_b, \quad \lambda_a^b(0) = \delta_a^b, \quad (30)$$

where we assume that

$$[\lambda_a^b, \lambda_c^d] = 0 \quad \Rightarrow \quad [\Omega_a, \Omega_b] = 0. \quad (31)$$

The quantum antibracket (16) is then given by

$$\begin{aligned} (\Omega_a, \Omega_b)_\Omega &= [\Omega_a, [\Omega, \Omega_b]] = -(i\hbar)^2 \lambda_a^f \lambda_b^e U_{ef}^d \mathcal{P}_d (-1)^{\varepsilon_d + \varepsilon_e + \varepsilon_f + \varepsilon_b \varepsilon_f} + \\ &+ i\hbar \left(\lambda_a^c [\theta'_c, \lambda_b^d] - \lambda_b^c [\theta'_c, \lambda_a^d] (-1)^{\varepsilon_a \varepsilon_b} \right) \mathcal{P}_d (-1)^{\varepsilon_c} - \\ &- i\hbar \left(\lambda_a^f \mathcal{C}^e [U_{ef}^c, \lambda_b^d] (-1)^{\varepsilon_b(\varepsilon_c+1)} - \lambda_b^f \mathcal{C}^e [U_{ef}^c, \lambda_a^d] (-1)^{\varepsilon_a(\varepsilon_c+1) + \varepsilon_a \varepsilon_b} \right) \mathcal{P}_d \mathcal{P}_c - \\ &- \mathcal{C}^e [[\theta'_e, \lambda_b^c], \lambda_a^d] \mathcal{P}_d \mathcal{P}_c (-1)^{(\varepsilon_a+1)(\varepsilon_b+\varepsilon_c+1)} - \\ &- \frac{1}{2} \mathcal{C}^f \mathcal{C}^e [[U_{ef}^c, \lambda_b^d], \lambda_a^g] \mathcal{P}_g \mathcal{P}_d \mathcal{P}_c (-1)^{\varepsilon_c + \varepsilon_f + (\varepsilon_a+1)(\varepsilon_b+\varepsilon_c+\varepsilon_d) + (\varepsilon_b+1)(\varepsilon_c+1)}. \end{aligned} \quad (32)$$

If we also require

$$(\Omega_a, \Omega_b, \Omega_c)_\Omega = 0 \quad \Leftrightarrow \quad [(\Omega_a, \Omega_b)_\Omega, \Omega_c] = 0, \quad (33)$$

then $(\Omega_a, \Omega_b)_\Omega$ in (32) satisfies the Jacobi identities which makes (15) integrable if $\Omega_{ab} = 0$. This condition is satisfied if we impose

$$[\lambda_a^b, U_{de}^c] = 0, \quad [\lambda_a^b, [\lambda_c^d, \theta_e]] = 0. \quad (34)$$

Eq.(15) may now be written as

$$\partial_a \tilde{\lambda}_b^c - \partial_b \tilde{\lambda}_a^c (-1)^{\varepsilon_a \varepsilon_b} = \tilde{\lambda}_a^e \tilde{\lambda}_b^d \tilde{U}_{de}^c (-1)^{\varepsilon_b \varepsilon_e + \varepsilon_c + \varepsilon_d + \varepsilon_e}. \quad (35)$$

where $\tilde{\lambda}_a^b \equiv V \lambda_a^b V^{-1}$ and $\tilde{U}_{ab}^c \equiv V U_{ab}^c V^{-1}$ where in turn the operator $V(\phi)$ is determined by the equation

$$i\hbar \partial_a V = V \lambda_a^b \theta'_b (-1)^{\varepsilon_b}. \quad (36)$$

Eq.(33) and $\Omega_{ab} = 0$ make all higher integrability conditions identically zero. One may note that

$$Y_a(\phi) = (i\hbar)^{-1} [\Omega, \Omega_a] = \lambda_a^b \theta'_b (-1)^{\varepsilon_a + \varepsilon_b} + \lambda_a^b \mathcal{C}^d U_{db}^c \mathcal{P}_c (-1)^{\varepsilon_a + \varepsilon_c} + (i\hbar)^{-1} \mathcal{C}^b [\theta_b, \lambda_a^c] \mathcal{P}_c. \quad (37)$$

5 Application: Generating operators for quantum antibrackets.

In [1] it was shown that quantum antibrackets for commuting operators or operators satisfying a nonabelian Lie algebra may be derived from a generating operator $Q(\phi)$. Here we generalize this construction to operators in arbitrary involutions. In distinction to the case in [1] we define here the generalized generating operator $Q(\phi)$ by equation (10), *i.e.*

$$Q(\phi) \overleftarrow{\nabla}_a \equiv Q(\phi) \overleftarrow{\partial}_a - (i\hbar)^{-1} [Q(\phi), Y_a(\phi)] = 0 \quad (38)$$

with the boundary condition

$$Q(0) = Q, \quad Q^2 = 0. \quad (39)$$

Since (38) implies

$$[Q(\phi), Q(\phi)] \overleftarrow{\partial}_a = -(i\hbar)^{-1} [Y_a(\phi), [Q(\phi), Q(\phi)]], \quad (40)$$

the boundary condition (39) implies $Q(\phi)^2 = 0$ which shows that $Q \rightarrow Q(\phi)$ is a unitary transformation.

Following ref.[1] we define generalized quantum antibrackets in terms of $Q(\phi)$ according to the formula

$$(Y_{a_1}, Y_{a_2}, \dots, Y_{a_n})'_{Q(\phi)} \equiv -Q(\phi) \overleftarrow{\partial}_{a_1} \overleftarrow{\partial}_{a_2} \cdots \overleftarrow{\partial}_{a_n} (i\hbar)^n (-1)^{E_n},$$

$$E_n \equiv \sum_{k=0}^{\lfloor \frac{n-1}{2} \rfloor} \varepsilon_{a_{2k+1}}. \quad (41)$$

In particular we have then (At $\phi^a = 0$ we have $Y_a(0) = \theta'_a + \{\text{possible ghost dependent terms}\}$.)

$$\begin{aligned} (Y_a(\phi), Y_b(\phi))'_{Q(\phi)} &\equiv -Q(\phi) \overleftarrow{\partial}_a \overleftarrow{\partial}_b (-1)^{\varepsilon_a} (i\hbar)^2 = \\ &= \frac{1}{2} \left([Y_a, [Q(\phi), Y_b]] - [Y_b, [Q(\phi), Y_a]] (-1)^{(\varepsilon_a+1)(\varepsilon_b+1)} \right) - \\ &\quad - \frac{1}{2} i\hbar [Q(\phi), Y_a \overleftarrow{\partial}_b + Y_b \overleftarrow{\partial}_a] (-1)^{\varepsilon_a \varepsilon_b} (-1)^{\varepsilon_a}, \end{aligned} \quad (42)$$

where we have made use of (10). The third line, which makes (42) differ from (16), is the price of a reparametrization independent extension of the definition (16) onto the space of parameters ϕ^a . The reason for this is that (42) is a second derivative of a scalar, which is not a tensor. Thus only within a preferred coordinate frame and at least at a fixed value of the parameters ϕ^a can one expect the formula (42) to reproduce the original antibracket (16). A similar situation is naturally expected to occur for all higher antibrackets as well. We also expect the canonical coordinates which are well known in Lie group theory, to be preferred ones also in the general case, so that all antibrackets would be reproduced in these coordinates at $\phi^a = 0$.

To illustrate the situation with the extension of the 3-antibracket, we give the following explicit formula

$$\begin{aligned}
& (Y_a(\phi), Y_b(\phi), Y_c(\phi))'_{Q(\phi)} (-1)^{(\varepsilon_a+1)(\varepsilon_c+1)} \equiv Q(\phi) \overleftarrow{\partial}_a \overleftarrow{\partial}_b \overleftarrow{\partial}_c (i\hbar)^3 (-1)^{\varepsilon_a \varepsilon_c} = \\
& = \frac{1}{3} \left([(Y_a, Y_b)'_Q, Y_c] (-1)^{\varepsilon_c + (\varepsilon_a+1)(\varepsilon_c+1)} + \text{cycle}(a, b, c) \right) + \\
& + \frac{1}{3} i\hbar \left([[Q, Y_a], Y_b \overleftarrow{\partial}_c + Y_c \overleftarrow{\partial}_b] (-1)^{\varepsilon_b \varepsilon_c} (-1)^{\varepsilon_a \varepsilon_c} + \text{cycle}(a, b, c) \right) + \\
& + \frac{1}{6} (i\hbar)^2 \left([Q, (Y_a \overleftarrow{\partial}_b + Y_b \overleftarrow{\partial}_a) (-1)^{\varepsilon_a \varepsilon_b}] \overleftarrow{\partial}_c (-1)^{\varepsilon_a \varepsilon_c} + \text{cycle}(a, b, c) \right). \tag{43}
\end{aligned}$$

Obviously the $\hbar$ -deviation from (25) vanishes together with the one for the 2-antibracket in (42). However, the vanishing of the $\hbar^2$ -deviation imposes a new condition. In the corresponding ϕ -extended n -antibrackets obtained from (41) we expect to have deviations involving up to $n - 2$ cyclically symmetrized derivatives of the 2-antibracket deviation which in terms of canonically coordinates should vanish at $\phi^a = 0$. The ϕ -extended n -antibrackets will then exactly reproduce the original n -antibrackets at $\phi^a = 0$.

For first rank theories equation (38) reduces to

$$Q(\phi) \overleftarrow{\partial}_a = (i\hbar)^{-1} [Q(\phi), \lambda_a^b(\phi) \theta'_b (-1)^{\varepsilon_a + \varepsilon_b}] \tag{44}$$

if we assume that $Q(\phi)$ commutes with the last terms in (37). Eq.(44) coincides exactly with equation (67) in appendix B of [1] if $U_{ab}^b(-1)^{\varepsilon_b} = 0$ or if $Q(\phi)$ commutes with $\lambda_a^b U_{bc}^c(-1)^{\varepsilon_c}$.

Acknowledgments

I.A.B. would like to thank Lars Brink for his very warm hospitality at the Institute of Theoretical Physics, Chalmers and Göteborg University. The work is partially supported by INTAS-RFBR grant 95-0829. The work of I.A.B. is also supported by INTAS grant 96-0308 and by RFBR grants 96-01-00482, 96-02-17314. I.A.B. and R.M. are thankful to the Royal Swedish Academy of Sciences for financial support.

References

- [1] I. A. Batalin and R. Marnelius, *Quantum antibrackets*, hep-th/9805084 (to be published in Phys. Lett. B).

- [2] I. A. Batalin and G. A. Vilkovisky, *Phys. Lett.* **B69**, 309 (1977),
E. S. Fradkin and T. E. Fradkina, *Phys. Lett.* **B72**, 343 (1978),
I. A. Batalin and E. S. Fradkin, *Phys. Lett.* **B122**, 157 (1983).
- [3] I. A. Batalin and E. S. Fradkin, *Phys. Lett.* **B128**, 303 (1983);
Riv. Nuovo Cim. **9**, 1 (1986); *Ann. Inst. Henri Poincaré* **49**, 145 (1988).
- [4] I. A. Batalin, *J. Math. Phys.* **22**, 1837 (1981)